

WARNING : Any malpractice or any attempt to commit any kind of malpractice in the Examination will **DISQUALIFY THE CANDIDATE.**

PAPER – II MATHEMATICS

Version Code	B1	Question Booklet Serial Number	
Time : 150 Minutes		Number of Questions : 120	Maximum Marks : 480
Name of Candidate			
Roll Number			
Signature of Candidate			

INSTRUCTIONS TO THE CANDIDATE

1. Please ensure that the **VERSION CODE** shown at the top of this Question Booklet is the same as that shown in the OMR Answer Sheet issued to you. If you have received a Question Booklet with a different **VERSION CODE**, please get it replaced with a Question Booklet with the same **VERSION CODE** as that of the OMR Answer Sheet from the Invigilator. **THIS IS VERY IMPORTANT.**
2. Please fill in the items such as name, signature and roll number of the candidate in the columns given above. Please also write the Question Booklet Sl. No. given at the top of this page against item 4 in the OMR Answer Sheet.
3. Please read the instructions given in the OMR Answer Sheet for marking answers. Candidates are advised to strictly follow the instructions contained in the OMR Answer Sheet.
4. This Question Booklet contains 120 Questions. For each Question, five answers are suggested and given against (A), (B), (C), (D) and (E) of which, only one will be the **Most Appropriate Answer**. Mark the bubble containing the letter corresponding to the 'Most Appropriate Answer' in the OMR Answer Sheet, by using either **Blue or Black ball - point pen only.**
5. Negative Marking: In order to discourage wild guessing, the score will be subject to penalization formula based on the number of right answers actually marked and the number of wrong answers marked. Each correct answer will be awarded FOUR marks. One mark will be deducted for each incorrect answer. More than one answer marked against a question will be deemed as incorrect answer and will be negatively marked.

IMMEDIATELY AFTER OPENING THIS QUESTION BOOKLET, THE CANDIDATE SHOULD VERIFY WHETHER THE QUESTION BOOKLET ISSUED CONTAINS ALL THE 120 QUESTIONS IN SERIAL ORDER. IF NOT, REQUEST FOR REPLACEMENT.

DO NOT OPEN THE SEAL UNTIL THE INVIGILATOR ASKS YOU TO DO SO.

Mathrubhumi Education

BLANK PAGE

Mathrubhumi Education



Maths-09/B1

2

**PLEASE ENSURE THAT THIS BOOKLET CONTAINS 120 QUESTIONS
SERIALLY NUMBERED FROM 1 TO 120 (Printed Pages : 32)**

1. The domain of the function $f(x) = \log_2(\log_3(\log_4 x))$ is
(A) $(-\infty, 4)$ (B) $(4, \infty)$ (C) $(0, 4)$ (D) $(1, \infty)$ (E) $(-\infty, 1)$
2. If $f: R \rightarrow R$ and $g: R \rightarrow R$ are defined by $f(x) = x - 3$ and $g(x) = x^2 + 1$, then the values of x for which $g(f(x)) = 10$ are
(A) $0, -6$ (B) $2, -2$ (C) $1, -1$ (D) $0, 6$ (E) $0, 2$
3. Two finite sets A and B have m and n elements respectively. If the total number of subsets of A is 112 more than the total number of subsets of B , then the value of m is
(A) 7 (B) 9 (C) 10 (D) 12 (E) 13
4. If $f(x)$ satisfies the relation $2f(x) + f(1-x) = x^2$ for all real x , then $f(x)$ is
(A) $\frac{x^2 + 2x - 1}{6}$ (B) $\frac{x^2 + 2x - 1}{3}$ (C) $\frac{x^2 + 4x - 1}{3}$
(D) $\frac{x^2 - 3x + 1}{6}$ (E) $\frac{x^2 + 3x - 1}{3}$

Space for rough work

5. The range of the function $f(x) = \frac{x^2 - x + 1}{x^2 + x + 1}$ where $x \in \mathbb{R}$, is
- (A) $(-\infty, 3]$ (B) $(-\infty, \infty)$ (C) $[3, \infty)$
(D) $[\frac{1}{3}, 3]$ (E) $(-\infty, \frac{1}{3}) \cup (3, \infty)$
6. If the area of the triangle formed by the points z , $z + iz$ and iz is 50 square units, then $|z|$ is equal to
- (A) 5 (B) 8 (C) 10 (D) 12 (E) $5\sqrt{2}$
7. The locus of z such that $\arg[(1 - 2i)z - 2 + 5i] = \frac{\pi}{4}$ is a
- (A) line not passing through the origin
(B) circle not passing through the origin
(C) line passing through the origin
(D) circle passing through the origin
(E) circle with centre at the origin
8. If $z = \sqrt{3} + i$, then the argument of $z^2 e^{z-i}$ is equal to
- (A) $\frac{\pi}{2}$ (B) $\frac{\pi}{6}$ (C) $e^{\frac{\pi}{6}}$ (D) $e^{\frac{\pi}{3}}$ (E) $\frac{\pi}{3}$

Space for rough work

Mathrubhumi
Education

9. If $\omega \neq 1$ and $\omega^3 = 1$, then $\frac{a\omega + b + c\omega^2}{a\omega^2 + b\omega + c} + \frac{a\omega^2 + b + c\omega}{a + b\omega + c\omega^2}$ is equal to
 (A) 2 (B) ω (C) 2ω (D) $2\omega^2$ (E) $a + b + c$
10. The centre of a regular hexagon is at the point $z = i$. If one of its vertices is at $2+i$, then the adjacent vertices of $2+i$ are at the points
 (A) $1 \pm 2i$ (B) $i + 1 \pm \sqrt{3}$ (C) $2 + i(1 \pm \sqrt{3})$
 (D) $1 + i(1 \pm \sqrt{3})$ (E) $1 - i(1 \pm \sqrt{3})$
11. If the roots of the equation $\frac{1}{x+p} + \frac{1}{x+q} = \frac{1}{r}$, ($x \neq -p, x \neq -q, r \neq 0$) are equal in magnitude but opposite in sign, then $p + q$ is equal to
 (A) r (B) $2r$ (C) r^2 (D) $\frac{1}{r}$ (E) $\frac{2}{r}$
12. The solution of the equation $(3 + 2\sqrt{2})^{x^2 - 8} + (3 + 2\sqrt{2})^{8 - x^2} = 6$ are
 (A) $3 \pm 2\sqrt{2}$ (B) ± 1 (C) $\pm 3\sqrt{3}, \pm 2\sqrt{2}$
 (D) $\pm 7, \pm \sqrt{3}$ (E) $\pm 3, \pm \sqrt{7}$
13. If $2 - i$ is a root of the equation $ax^2 + 12x + b = 0$ (where a and b are real), then the value of ab is equal to
 (A) 45 (B) 15 (C) -15 (D) -45 (E) 25

Space for rough work

14. If one root of the equation $lx^2 + mx + n = 0$ is $\frac{9}{2}$ (l, m and n are positive integers) and $\frac{m}{4n} = \frac{l}{m}$, then $l + n$ is equal to
 (A) 80 (B) 85 (C) 90 (D) 95 (E) 100
15. If $x^2 + 4ax + 2 > 0$ for all values of x , then a lies in the interval
 (A) $(-2, 4)$ (B) $(1, 2)$ (C) $(-\sqrt{2}, \sqrt{2})$
 (D) $(-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}})$ (E) $(-4, 2)$
16. If a, b, c are in G.P. and x, y are arithmetic mean of a, b and b, c respectively, then $\frac{1}{x} + \frac{1}{y}$ is equal to
 (A) $\frac{2}{b}$ (B) $\frac{3}{b}$ (C) $\frac{b}{3}$ (D) $\frac{b}{2}$ (E) $\frac{1}{b}$
17. A student read common difference of an A.P. as -3 instead of 3 and obtained the sum of first 10 terms as -30 . Then the actual sum of first 10 terms is equal to
 (A) 240 (B) 120 (C) 300 (D) 180 (E) 480
18. If $a_1 = 1$ and $a_n = n a_{n-1}$ for all positive integer $n \geq 2$ then a_5 is equal to
 (A) 125 (B) 120 (C) 100 (D) 24 (E) 6

Space for rough work

19. If a_1, a_2, \dots, a_n are in A.P. with common difference $d \neq 0$, then $(\sin d)[\sec a_1 \sec a_2 + \sec a_2 \sec a_3 + \dots + \sec a_{n-1} \sec a_n]$ is equal to
- (A) $\cot a_n - \cot a_1$ (B) $\cot a_1 - \cot a_n$ (C) $\tan a_n - \tan a_1$
(D) $\tan a_n - \tan a_{n-1}$ (E) $\tan a_1 - \tan a_n$
20. The value of $\frac{1}{2!} + \frac{2}{3!} + \dots + \frac{999}{1000!}$ is equal to
- (A) $\frac{1000! - 1}{1000!}$ (B) $\frac{1000! + 1}{1000!}$ (C) $\frac{999! - 1}{999!}$
(D) $\frac{999! + 1}{999!}$ (E) $\frac{1000! - 999!}{1000!}$
21. $\log_e \frac{1+3x}{1-2x}$ is equal to
- (A) $-5x - \frac{5x^2}{2} - \frac{35x^3}{3} - \dots$ (B) $-5x + \frac{5x^2}{2} - \frac{35x^3}{3} + \dots$ (C) $5x - \frac{5x^2}{2} + \frac{35x^3}{3} - \dots$
(D) $5x + \frac{5x^2}{2} + \frac{35x^3}{3} + \dots$ (E) $x + \frac{3x^2}{2} + \frac{5x^3}{4} + \dots$
22. The sum of the infinite series $\frac{1}{2}\left(\frac{1}{3} + \frac{1}{4}\right) - \frac{1}{4}\left(\frac{1}{3^2} + \frac{1}{4^2}\right) + \frac{1}{6}\left(\frac{1}{3^3} + \frac{1}{4^3}\right) - \dots$ is equal to
- (A) $\frac{1}{2} \log 2$ (B) $\log \frac{3}{5}$ (C) $\log \frac{5}{3}$ (D) $\frac{1}{2} \log \frac{5}{3}$ (E) $\frac{1}{2} \log \frac{3}{5}$
23. The sum of the infinite series $\frac{2^2}{2!} + \frac{2^4}{4!} + \frac{2^6}{6!} + \dots$ is equal to
- (A) $\frac{e^2 + 1}{2e}$ (B) $\frac{e^4 + 1}{2e^2}$ (C) $\frac{(e^2 - 1)^2}{2e^2}$ (D) $\frac{(e^2 + 1)^2}{2e^2}$ (E) $\frac{(e^2 - 1)^2}{4e^2}$

Space for rough work

24. If $|x| < 1$, then the coefficient of x^6 in the expansion of $(1+x+x^2)^{-3}$ is
 (A) 3 (B) 6 (C) 9 (D) 12 (E) 15
25. ${}^{15}C_0 \cdot {}^5C_5 + {}^{15}C_1 \cdot {}^5C_4 + {}^{15}C_2 \cdot {}^5C_3 + {}^{15}C_3 \cdot {}^5C_2 + {}^{15}C_4 \cdot {}^5C_1$ is equal to
 (A) $2^{20} - 2^5$ (B) $\frac{20!}{5!15!}$ (C) $\frac{20!}{5!15!} - 1$
 (D) $\frac{20!}{5!15!} - \frac{15!}{5!10!}$ (E) $\frac{15!}{5!10!}$
26. If ${}^{2n+1}P_{n-1} : {}^{2n-1}P_n = 3 : 5$ then the value of n is equal to
 (A) 4 (B) 3 (C) 2 (D) 1 (E) 5
27. Let $[x]$ denote the greatest integer less than or equal to x . If $x = (\sqrt{3}+1)^5$, then $[x]$ is equal to
 (A) 75 (B) 50 (C) 76 (D) 51 (E) 152
28. If n is a positive integer, then $5^{2n+2} - 24n - 25$ is divisible by
 (A) 574 (B) 575 (C) 675 (D) 674 (E) 576
29. Let T_n denote the number of triangles which can be formed by using the vertices of a regular polygon of n sides. If $T_{n+1} - T_n = 28$, then n equals
 (A) 4 (B) 5 (C) 6 (D) 7 (E) 8

Space for rough work

30. If α, β, γ are the cube roots of unity then the value of the determinant

$$\begin{vmatrix} e^\alpha & e^{2\alpha} & (e^{3\alpha} - 1) \\ e^\beta & e^{2\beta} & (e^{3\beta} - 1) \\ e^\gamma & e^{2\gamma} & (e^{3\gamma} - 1) \end{vmatrix} \text{ is equal to}$$

- (A) -2 (B) -1 (C) 0 (D) 1 (E) 2

31. If B is a non-singular matrix and A is a square matrix such that $B^{-1}AB$ exists, then $\det(B^{-1}AB)$ is equal to

- (A) $\det(A^{-1})$ (B) $\det(B^{-1})$ (C) $\det(B)$
 (D) $\det(A)$ (E) $\det(AB^{-1})$

32. If $1, \omega, \omega^2$ are cube roots of unity and if $\begin{bmatrix} 1+\omega & 2\omega \\ -2\omega & -b \end{bmatrix} + \begin{bmatrix} a & -\omega \\ 3\omega & 2 \end{bmatrix} = \begin{bmatrix} 0 & \omega \\ \omega & 1 \end{bmatrix}$, then $a^2 + b^2$ is equal to

- (A) $1 + \omega^2$ (B) $\omega^2 - 1$ (C) $1 + \omega$ (D) $(1 + \omega)^2$ (E) ω^2

33. If the three linear equations

$$x + 4ay + az = 0$$

$$x + 3by + bz = 0$$

$$x + 2cy + cz = 0$$

have a non-trivial solution, where $a \neq 0, b \neq 0, c \neq 0$, then $ab + bc$ is equal to

- (A) $2ac$ (B) $-ac$ (C) ac (D) $-2ac$ (E) a

34. If $A = \begin{bmatrix} 1 & 0 & 0 \\ x & 1 & 0 \\ x & x & 1 \end{bmatrix}$ and $I = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ then $A^3 - 3A^2 + 3A$ is equal to


- (A) $3I$ (B) I (C) $-I$ (D) $-2I$ (E) $2I$

Space for rough work

35. If $A = \begin{pmatrix} 1 & 2 \\ 3 & 5 \end{pmatrix}$, then the value of the determinant $|A^{2009} - 5A^{2008}|$ is
 (A) -6 (B) -5 (C) -4 (D) 4 (E) 6
36. If x satisfies the inequations $2x - 7 < 11$, $3x + 4 < -5$, then x lies in the interval
 (A) $(-\infty, 3)$ (B) $(-\infty, 2)$ (C) $(-\infty, -3)$ (D) $(-\infty, \infty)$ (E) $(3, \infty)$
37. The set of all real x satisfying the inequality $\frac{3-|x|}{4-|x|} \geq 0$ is
 (A) $[-3, 3] \cup (-\infty, -4) \cup (4, \infty)$ (B) $(-\infty, -4) \cup (4, \infty)$
 (C) $(-\infty, -3) \cup (4, \infty)$ (D) $(-\infty, -3) \cup (3, \infty)$
 (E) $[-3, 3] \cup [4, \infty)$
38. Identify the false statement
 (A) $\sim [p \vee (\sim q)] \equiv (\sim p) \wedge q$
 (B) $[p \vee q] \vee (\sim p)$ is a tautology
 (C) $[p \wedge q] \wedge (\sim p)$ is a contradiction
 (D) $\sim [p \wedge (\sim p)]$ is a tautology
 (E) $\sim (p \vee q) \equiv (\sim p) \vee (\sim q)$

Space for rough work

39. The Boolean expression corresponding to the combinational circuit is



- (A) $(x_1 + x_2 \cdot x_3') \cdot x_2$ (B) $(x_1 \cdot (x_2 + x_3)) + x_2$
 (C) $(x_1 \cdot (x_2 + x_3')) + x_2$ (D) $(x_1 \cdot (x_2 + x_3')) + x_3$
 (E) $(x_1 + x_2' \cdot x_3) \cdot x_2$

40. In a Boolean algebra B with respect to '+' and ' \cdot ', x' denotes the negation of $x \in B$. Then

- (A) $x - x' = 1$ and $x \cdot x' = 1$ (B) $x + x' = 1$ and $x \cdot x' = 0$
 (C) $x + x' = 0$ and $x \cdot x' = 1$ (D) $x + x' = 0$ and $x \cdot x' = 0$
 (E) $x - x' = 0$ and $x \cdot x' = 0$

41. If $\cos^{-1}\left(\frac{5}{13}\right) - \sin^{-1}\left(\frac{12}{13}\right) = \cos^{-1} x$, then x is equal to

- (A) 1 (B) $\frac{1}{\sqrt{2}}$ (C) 0 (D) $\frac{\sqrt{3}}{2}$ (E) -1

Space for rough work

42. The value of $\cos(\tan^{-1}(\sin(\cot^{-1}x)))$ is
- (A) $\sqrt{\frac{x^2+1}{x^2-1}}$ (B) $\sqrt{\frac{1-x^2}{x^2+2}}$ (C) $\sqrt{\frac{1-x^2}{1+x^2}}$ (D) $\sqrt{\frac{x^2+1}{x^2+2}}$ (E) $\sqrt{\frac{1-x^2}{2-x^2}}$
43. If a and b are positive numbers such that $a > b$, then the minimum value of $a \sec\theta - b \tan\theta$ ($0 < \theta < \frac{\pi}{2}$) is
- (A) $\frac{1}{\sqrt{a^2-b^2}}$ (B) $\frac{1}{\sqrt{a^2+b^2}}$ (C) $\sqrt{a^2+b^2}$ (D) $\sqrt{a^2-b^2}$ (E) a^2-b^2
44. If $-\frac{\pi}{2} < \sin^{-1}x < \frac{\pi}{2}$, then $\tan(\sin^{-1}x)$ is equal to
- (A) $\frac{x}{1-x^2}$ (B) $\frac{x}{1+x^2}$ (C) $\frac{x}{\sqrt{1-x^2}}$ (D) $\frac{1}{\sqrt{1-x^2}}$ (E) $\frac{x}{\sqrt{x^2-1}}$
45. If $A+B=45^\circ$ then $(\cot A-1)(\cot B-1)$ is equal to
- (A) 1 (B) $\frac{1}{2}$ (C) -1 (D) -2 (E) 2
46. The solution of the equation $[\sin x + \cos x]^{1+\sin 2x} = 2$, $-\pi \leq x \leq \pi$ is
- (A) $\frac{\pi}{2}$ (B) π (C) $\frac{\pi}{4}$ (D) $\frac{3\pi}{4}$ (E) $\frac{\pi}{3}$
47. If $\sin A - \sqrt{6} \cos A = \sqrt{7} \cos A$, then $\cos A + \sqrt{6} \sin A$ is equal to
- (A) $\sqrt{6} \sin A$ (B) $\sqrt{7} \sin A$ (C) $\sqrt{6} \cos A$ (D) $\sqrt{7} \cos A$ (E) $\sqrt{42} \cos A$

Space for rough work

48. If $\tan A$ and $\tan B$ are the roots of $abx^2 - c^2x + ab = 0$ where a, b, c are the sides of the triangle ABC , then the value of $\sin^2 A + \sin^2 B + \sin^2 C$ is
 (A) 1 (B) 3 (C) 4 (D) 2 (E) 5
49. In a triangle ABC , if $a = 3, b = 4, c = 5$ then the distance between its incentre and circumcentre is
 (A) $\frac{1}{2}$ (B) $\frac{\sqrt{3}}{2}$ (C) $\frac{3}{2}$ (D) $\frac{5}{2}$ (E) $\frac{\sqrt{5}}{2}$
50. In triangle ABC , the value of $\frac{\cot \frac{A}{2} \cot \frac{B}{2} - 1}{\cot \frac{A}{2} \cot \frac{B}{2}}$ is
 (A) $\frac{a}{a+b+c}$ (B) $\frac{c}{a+b+c}$ (C) $\frac{2a}{a+b+c}$ (D) $\frac{2b}{a+b+c}$ (E) $\frac{2c}{a+b+c}$
51. In a triangle ABC if $\angle A = 60^\circ, a = 5, b = 4$, then c is a root of the equation
 (A) $c^2 - 5c - 9 = 0$ (B) $c^2 - 4c - 9 = 0$ (C) $c^2 - 10c + 25 = 0$
 (D) $c^2 - 5c - 41 = 0$ (E) $c^2 - 4c - 41 = 0$
52. From the top of a tower, the angle of depression of a point on the ground is 60° . If the distance of this point from the tower is $\frac{1}{\sqrt{3}+1}$ metres, then the height of the tower is
 (A) $\frac{4\sqrt{3}}{2}$ metres (B) $\frac{\sqrt{3}+3}{2}$ metres (C) $\frac{3-\sqrt{3}}{2}$ metres
 (D) $\frac{\sqrt{3}}{2}$ metres (E) $\sqrt{3}+1$ metres

Space for rough work

53. The vertices of a family of triangles have integer co-ordinates. If two of the vertices of all the triangles are (0, 0) and (6, 8), then the least value of areas of the triangles is
- (A) 1 (B) $\frac{3}{2}$ (C) 2 (D) $\frac{5}{2}$ (E) 3
54. A line has slope m and y -intercept 4. The distance between the origin and the line is equal to
- (A) $\frac{4}{\sqrt{1-m^2}}$ (B) $\frac{4}{\sqrt{m^2-1}}$ (C) $\frac{4}{\sqrt{m^2+1}}$ (D) $\frac{4m}{\sqrt{1+m^2}}$ (E) $\frac{4m}{\sqrt{m^2-1}}$
55. One side of length $3a$ of a triangle of area a^2 square units lies on the line $x = a$. Then one of the lines on which the third vertex lies, is
- (A) $x = -a^2$ (B) $x = a^2$ (C) $x = -a$ (D) $x = \frac{a}{3}$ (E) $x = -\frac{a}{3}$
56. The distance of the point (1, 2) from the line $x + y + 5 = 0$ measured along the line parallel to $3x - y = 7$ is equal to
- (A) $4\sqrt{10}$ (B) 40 (C) $\sqrt{40}$ (D) $10\sqrt{2}$ (E) $2\sqrt{20}$
57. Area of the triangle formed by the lines $y = 2x$, $y = 3x$ and $y = 5$ is equal to (in square units)
- (A) $\frac{25}{6}$ (B) $\frac{25}{12}$ (C) $\frac{5}{6}$ (D) $\frac{17}{12}$ (E) 6
58. Triangle ABC has vertices (0, 0), (11, 60) and (91, 0). If the line $y = kx$ cuts the triangle into two triangles of equal area, then k is equal to
- (A) $\frac{30}{51}$ (B) $\frac{4}{7}$ (C) $\frac{7}{4}$ (D) $\frac{30}{91}$ (E) $\frac{25}{37}$

Space for rough work

59. If the lines $y = 3x + 1$ and $2y = x + 3$ are equally inclined to the line $y = mx + 4$, $\left(\frac{1}{2} < m < 3\right)$, then the values of m are
- (A) $\frac{1}{7}(1 \pm 5\sqrt{3})$ (B) $\frac{1}{7}(1 \pm 5\sqrt{5})$ (C) $\frac{1}{7}(1 \pm 5\sqrt{2})$
(D) $\frac{1}{7}(1 \pm 2\sqrt{5})$ (E) $\frac{1}{7}(1 \pm 3\sqrt{2})$
60. The vertices of a triangle are $(3, 0)$, $(3, 3)$ and $(0, 3)$. Then the coordinates of the circumcentre are
- (A) $(0, 0)$ (B) $(1, 1)$ (C) $\left(\frac{5}{2}, \frac{5}{2}\right)$ (D) $(2, 2)$ (E) $\left(\frac{3}{2}, \frac{3}{2}\right)$
61. Area of the equilateral triangle inscribed in the circle $x^2 + y^2 - 7x + 9y + 5 = 0$ is
- (A) $\frac{155}{8}\sqrt{3}$ square units (B) $\frac{165}{8}\sqrt{3}$ square units (C) $\frac{175}{8}\sqrt{3}$ square units
(D) $\frac{185}{8}\sqrt{3}$ square units (E) $\frac{195}{8}\sqrt{3}$ square units
62. The equation of one of the diameters of the circle $x^2 + y^2 - 6x + 2y = 0$ is
- (A) $x + y = 0$ (B) $x - y = 0$ (C) $3x + y = 0$ (D) $x + 3y = 0$ (E) $x + 2y = 0$
63. If two chords having lengths $a^2 - 1$ and $3(a + 1)$, where a is a constant, of a circle bisect each other, then the radius of the circle is
- (A) 6 (B) $\frac{15}{2}$ (C) 8 (D) $\frac{19}{2}$ (E) 10

Space for rough work

64. The equation of the parabola whose focus (3, 2) and vertex (1, 2), is
- (A) $x^2 + 4x - 8y + 12 = 0$ (B) $x^2 - 4x - 8y + 12 = 0$
 (C) $y^2 - 8x - 4y + 12 = 0$ (D) $y^2 + 4y - 8x + 12 = 0$
 (E) $y^2 - 8x - 2y - 17 = 0$
65. The sum of the distances of a point (2, -3) from the foci of an ellipse $16(x-2)^2 + 25(y+3)^2 = 400$ is
- (A) 8 (B) 6 (C) 50 (D) 32 (E) 10
66. The equation of one of the tangents to $\frac{x^2}{3} - \frac{y^2}{2} = 1$ which is parallel to $y = x$, is
- (A) $x - y + 2 = 0$ (B) $x + y - 1 = 0$ (C) $x + y - 2 = 0$
 (D) $x - y + 1 = 0$ (E) $x + y + 1 = 0$
67. If e_1 is the eccentricity of the ellipse $\frac{x^2}{16} + \frac{y^2}{7} = 1$ and e_2 is the eccentricity of the hyperbola $\frac{x^2}{9} - \frac{y^2}{7} = 1$, then $e_1 + e_2$ is equal to
- (A) $\frac{16}{7}$ (B) $\frac{25}{4}$ (C) $\frac{25}{12}$ (D) $\frac{16}{9}$ (E) $\frac{23}{16}$
68. If $\vec{p} \times \vec{q} = \vec{r}$ and $\vec{q} \times \vec{r} = \vec{p}$, then
- (A) $r = 1, p = q$ (B) $p = 1, q = 1$ (C) $r = 2p, q = 2$
 (D) $q = 1, p = r$ (E) $q = 1, r = 1$

Space for rough work

69. Vectors \vec{a} and \vec{b} are inclined at an angle $\theta = 120^\circ$. If $|\vec{a}|=1, |\vec{b}|=2$ then $[(\vec{a} + 3\vec{b}) \times (3\vec{a} + \vec{b})]^2$ is equal to
 (A) 190 (B) 275 (C) 300 (D) 320 (E) 192
70. If the projection of the vector \vec{a} on \vec{b} is $|\vec{a} \times \vec{b}|$ and if $3\vec{b} = \vec{i} + \vec{j} + \vec{k}$, then the angle between \vec{a} and \vec{b} is
 (A) $\pi/3$ (B) $\pi/2$ (C) $\pi/4$ (D) $\pi/6$ (E) 0
71. If $\vec{x} = \vec{a} + \vec{b}, \vec{y} = \vec{a} - \vec{b}, |\vec{a}| = 2, |\vec{b}| = 3$ and the angle between \vec{a} and \vec{b} is $\frac{\pi}{3}$, then $|\vec{x} \times \vec{y}|$ is equal to
 (A) $5\sqrt{3}$ (B) 6 (C) $4\sqrt{3}$ (D) 9 (E) $6\sqrt{3}$
72. If the position vectors of three consecutive vertices of a parallelogram are $\vec{i} + \vec{j} + \vec{k}, \vec{i} + 3\vec{j} + 5\vec{k}$ and $7\vec{i} + 9\vec{j} + 11\vec{k}$, then the coordinates of the fourth vertex are
 (A) (2, 1, 3) (B) (6, 7, 8) (C) (4, 1, 3) (D) (7, 7, 7) (E) (8, 8, 8)
73. The two variable vectors $3x\vec{i} + y\vec{j} - 3\vec{k}$ and $x\vec{i} - 4y\vec{j} + 4\vec{k}$ are orthogonal to each other, then the locus of (x, y) is
 (A) hyperbola (B) circle (C) straight line
 (D) ellipse (E) parabola

Space for rough work

74. If $\vec{a}, \vec{b}, \vec{c}$ are non-coplanar and $(\vec{a} + \lambda\vec{b}) \cdot [(\vec{b} + 3\vec{c}) \times (\vec{c} - 4\vec{a})] = 0$, then the value of λ is equal to
- (A) 0 (B) $\frac{1}{12}$ (C) $\frac{5}{12}$ (D) 3 (E) $\frac{7}{12}$
75. The angle between the line $\frac{3x-1}{3} = \frac{y+3}{-1} = \frac{5-2z}{4}$ and the plane $3x-3y-6z = 10$ is equal to
- (A) $\frac{\pi}{6}$ (B) $\frac{\pi}{4}$ (C) $\frac{\pi}{3}$ (D) $\frac{\pi}{2}$ (E) $\frac{2\pi}{3}$
76. The angle between the straight lines $\vec{r} = (2-3t)\vec{i} + (1+2t)\vec{j} + (2+6t)\vec{k}$ and $\vec{r} = (1+4s)\vec{i} + (2-s)\vec{j} + (8s-1)\vec{k}$ is
- (A) $\cos^{-1}\left(\frac{\sqrt{41}}{34}\right)$ (B) $\cos^{-1}\left(\frac{21}{34}\right)$ (C) $\cos^{-1}\left(\frac{43}{63}\right)$
(D) $\cos^{-1}\left(\frac{5\sqrt{23}}{41}\right)$ (E) $\cos^{-1}\left(\frac{34}{63}\right)$
77. If Q is the image of the point $P(2, 3, 4)$ under the reflection in the plane $x - 2y + 5z = 6$, then the equation of the line PQ is
- (A) $\frac{x-2}{-1} = \frac{y-3}{2} = \frac{z-4}{5}$ (B) $\frac{x-2}{1} = \frac{y-3}{-2} = \frac{z-4}{5}$
(C) $\frac{x-2}{-1} = \frac{y-3}{-2} = \frac{z-4}{5}$ (D) $\frac{x-2}{1} = \frac{y-3}{2} = \frac{z-4}{5}$
(E) $\frac{x-2}{1} = \frac{y-3}{2} = \frac{z+4}{5}$

Space for rough work

78. The distance of the point of intersection of the line $\frac{x-2}{3} = \frac{y+1}{4} = \frac{z-2}{12}$ and the plane $x - y + z = 5$ from the point $(-1, -5, -10)$ is
 (A) 13 (B) 12 (C) 11 (D) 8 (E) 7
79. If the direction cosines of a line are $\left(\frac{1}{c}, \frac{1}{c}, \frac{1}{c}\right)$, then
 (A) $0 < c < 1$ (B) $c > 2$ (C) $c = \pm\sqrt{2}$ (D) $c = \pm\sqrt{3}$ (E) $c = \pm 3$
80. The vector form of the sphere $2(x^2 + y^2 + z^2) - 4x + 6y + 8z - 5 = 0$ is
 (A) $\vec{r} \cdot [\vec{r} - (2\vec{i} + \vec{j} + \vec{k})] = \frac{2}{5}$ (B) $\vec{r} \cdot [\vec{r} - (2\vec{i} - 3\vec{j} - 4\vec{k})] = \frac{1}{2}$
 (C) $\vec{r} \cdot [\vec{r} - (2\vec{i} + 3\vec{j} + 4\vec{k})] = \frac{5}{2}$ (D) $\vec{r} \cdot [\vec{r} + (2\vec{i} - 3\vec{j} - 4\vec{k})] = \frac{5}{2}$
 (E) $\vec{r} \cdot [\vec{r} - (2\vec{i} - 3\vec{j} - 4\vec{k})] = \frac{5}{2}$
81. If the lines $\frac{1-x}{3} = \frac{y-2}{2\alpha} = \frac{z-3}{2}$ and $\frac{x-1}{3\alpha} = y-1 = \frac{6-z}{5}$ are perpendicular, then the value of α is
 (A) $\frac{-10}{7}$ (B) $\frac{10}{7}$ (C) $\frac{-10}{11}$ (D) $\frac{10}{11}$ (E) $\frac{10}{9}$
82. The distance between the lines $\vec{r} = (4\vec{i} - 7\vec{j} - 9\vec{k}) + t(3\vec{i} - 7\vec{j} + 4\vec{k})$ and $\vec{r} = (7\vec{i} - 14\vec{j} - 5\vec{k}) + s(-3\vec{i} + 7\vec{j} - 4\vec{k})$ is equal to
 (A) 1 (B) $\frac{1}{2}$ (C) $\frac{3}{4}$ (D) 17 (E) 0

Space for rough work

83. If the variance of $1, 2, 3, 4, 5, \dots, 10$ is $\frac{99}{12}$, then the standard deviation of $3, 6, 9, 12, \dots, 30$ is
- (A) $\frac{297}{4}$ (B) $\frac{3}{2}\sqrt{33}$ (C) $\frac{3}{2}\sqrt{99}$ (D) $\sqrt{\frac{99}{12}}$ (E) $\frac{3\sqrt{3}}{2}$
84. The mean of the values $0, 1, 2, 3, \dots, n$ with the corresponding weights ${}^n C_0, {}^n C_1, \dots, {}^n C_n$ respectively, is
- (A) $\frac{n+1}{2}$ (B) $\frac{n-1}{2}$ (C) $\frac{2^n-1}{2}$ (D) $\frac{2^n+1}{2}$ (E) $\frac{n}{2}$
85. A complete cycle of a traffic light takes 60 seconds. During each cycle the light is green for 25 seconds, yellow for 5 seconds and red for 30 seconds. At a randomly chosen time, the probability that the light will not be green is
- (A) $\frac{1}{3}$ (B) $\frac{1}{4}$ (C) $\frac{4}{12}$ (D) $\frac{7}{12}$ (E) $\frac{3}{4}$
86. If the random variable X takes the values $x_1, x_2, x_3, \dots, x_{10}$ with probabilities $P(X = x_i) = ki$, then the value of k is equal to
- (A) $\frac{1}{10}$ (B) $\frac{1}{15}$ (C) $\frac{1}{55}$ (D) 10 (E) 55
87. Let α and β be the roots of $ax^2 + bx + c = 0$. Then $\lim_{x \rightarrow \alpha} \frac{1 - \cos(ax^2 + bx + c)}{(x - \alpha)^2}$ is equal to
- (A) 0 (B) $\frac{1}{2}(\alpha - \beta)^2$ (C) $\frac{a^2}{2}(\alpha - \beta)^2$ (D) $(\alpha - \beta)$ (E) 1

Space for rough work

88. The number of discontinuities of the greatest integer function $f(x) = [x]$, $x \in \left(-\frac{7}{2}, 100\right)$ is equal to
 (A) 104 (B) 100 (C) 102 (D) 101 (E) 103
89. If $f(x) = \begin{cases} \frac{3 \sin \pi x}{5x}, & x \neq 0 \\ 2k, & x = 0 \end{cases}$ is continuous at $x = 0$, then the value of k is equal to
 (A) $\frac{3\pi}{10}$ (B) $\frac{3\pi}{5}$ (C) $\frac{\pi}{10}$ (D) $\frac{3\pi}{2}$ (E) $\frac{2\pi}{3}$
90. If a function f satisfies $f(f(x)) = x + 1$ for all real values of x and if $f(0) = \frac{1}{2}$, then $f(1)$ is equal to
 (A) $\frac{1}{2}$ (B) 1 (C) $\frac{3}{2}$ (D) 2 (E) 0
91. If $y = \log_2 \log_2(x)$ then $\frac{dy}{dx} =$
 (A) $\frac{\log_2 e}{\log_e x}$ (B) $\frac{\log_2 e}{x \log_x 2}$ (C) $\frac{\log_2 x}{\log_e 2}$ (D) $\frac{\log_2 e}{\log_2 x}$ (E) $\frac{\log_2 e}{x \log_e x}$
92. If $\frac{d}{dx}(f(x)) = \frac{1}{1+x^2}$ then $\frac{d}{dx}(f(x^3))$ is
 (A) $\frac{3x}{1+x^3}$ (B) $\frac{3x^2}{1+x^6}$ (C) $\frac{-6x^5}{(1+x^6)^2}$ (D) $\frac{-6x^5}{1+x^6}$ (E) $\tan^{-1} x$

Space for rough work

93. If $y = \sin\{\cos^{-1}[\sin(\cos^{-1} x)]\}$, then $\frac{dy}{dx}$ at $x = \frac{1}{2}$ is equal to
 (A) 0 (B) -1 (C) $\frac{2}{\sqrt{3}}$ (D) $\frac{1}{\sqrt{3}}$ (E) 1
94. If $x^2 + y^2 = t - \frac{1}{t}$ and $x^4 + y^4 = t^2 + \frac{1}{t^2}$ then $\frac{dy}{dx}$ is equal to
 (A) $\frac{1}{x^2 y^3}$ (B) $\frac{1}{xy^3}$ (C) $\frac{1}{x^2 y^2}$ (D) $\frac{1}{x^3 y}$ (E) $\frac{-1}{x^3 y}$
95. If $y = \sec^{-1}[\operatorname{cosec} x] + \operatorname{cosec}^{-1}[\sec x] + \sin^{-1}[\cos x] + \cos^{-1}[\sin x]$, then $\frac{dy}{dx}$ is equal to
 (A) 0 (B) 2 (C) -2 (D) -4 (E) 1
96. If $y = e^x \cdot e^{x^2} \cdot e^{x^3} \dots e^{x^n} \dots$, for $0 < x < 1$, then $\frac{dy}{dx}$ at $x = \frac{1}{2}$ is
 (A) e (B) $4e$ (C) $2e$ (D) $3e$ (E) $5e$
97. The derivative of $\tan^{-1}\left(\frac{2x}{1-x^2}\right)$ with respect to $\cos^{-1}\sqrt{1-x^2}$ is
 (A) $\frac{\sqrt{1-x^2}}{1+x^2}$ (B) $\frac{1}{\sqrt{1-x^2}}$ (C) $\frac{2}{\sqrt{1-x^2}(1+x^2)}$
 (D) $\frac{2}{1+x^2}$ (E) $\frac{2\sqrt{1-x^2}}{1+x^2}$

Space for rough work

98. If the curves $\frac{x^2}{a^2} + \frac{y^2}{12} = 1$ and $y^3 = 8x$ intersect at right angles, then the value of a^2 is equal to
(A) 16 (B) 12 (C) 8 (D) 4 (E) 2
99. If the function $f(x) = x^3 - 12ax^2 + 36a^2x - 4$ ($a > 0$) attains its maximum and minimum at $x = p$ and $x = q$ respectively and if $3p = q^2$, then a is equal to
(A) $\frac{1}{6}$ (B) $\frac{1}{36}$ (C) $\frac{1}{3}$ (D) 18 (E) 6
100. The equation of the tangent to the curve $y = 4e^{-\frac{x}{4}}$ at the point where the curve crosses y -axis is equal to
(A) $3x + 4y = 16$ (B) $4x + y = 4$ (C) $x + y = 4$
(D) $4x - 3y = -12$ (E) $x - y = -4$
101. The diagonal of a square is changing at the rate of 0.5 cm/sec. Then the rate of change of area, when the area is 400 cm^2 , is equal to
(A) $20\sqrt{2} \text{ cm}^2/\text{sec}$ (B) $10\sqrt{2} \text{ cm}^2/\text{sec}$ (C) $\frac{1}{10\sqrt{2}} \text{ cm}^2/\text{sec}$
(D) $\frac{10}{\sqrt{2}} \text{ cm}^2/\text{sec}$ (E) $5\sqrt{2} \text{ cm}^2/\text{sec}$

Space for rough work

102. The equation of the tangent to the curve $x^2 - 2xy + y^2 + 2x + y - 6 = 0$ at $(2, 2)$ is

- (A) $2x + y - 6 = 0$ (B) $2y + x - 6 = 0$ (C) $x + 3y - 8 = 0$
(D) $3x + y - 8 = 0$ (E) $x + y - 4 = 0$

103. The angle between the curves $y = a^x$ and $y = b^x$ is equal to

- (A) $\tan^{-1}\left(\frac{a-b}{1+ab}\right)$ (B) $\tan^{-1}\left(\frac{a+b}{1-ab}\right)$
(C) $\tan^{-1}\left(\frac{\log b + \log a}{1 + \log a \log b}\right)$ (D) $\tan^{-1}\left(\frac{\log a + \log b}{1 - \log a \log b}\right)$
(E) $\tan^{-1}\left(\frac{\log a - \log b}{1 + \log a \log b}\right)$

104. Let $f(x) = (x-7)^2(x-2)^7$, $x \in [2, 7]$. The value of $\theta \in (2, 7)$ such that $f'(\theta) = 0$ is equal to

- (A) $\frac{49}{4}$ (B) $\frac{53}{9}$ (C) $\frac{53}{7}$ (D) $\frac{49}{9}$ (E) $\frac{45}{7}$

Space for rough work

105. $\int (\sqrt[3]{x}) \left(\sqrt[5]{1+\sqrt[3]{x^4}} \right) dx$ is equal to

- (A) $\left(1+x^{\frac{3}{4}}\right)^{\frac{6}{5}} + C$ (B) $\left(1+x^{\frac{4}{3}}\right)^{\frac{6}{5}} + C$ (C) $\frac{5}{8} \left(1+x^{\frac{4}{3}}\right)^{\frac{6}{5}} + C$
(D) $\frac{1}{6} \left(1+x^{\frac{4}{3}}\right)^{\frac{6}{5}} + C$ (E) $\frac{15}{8} \left(1+x^{\frac{4}{3}}\right)^{\frac{6}{5}} + C$

106. If $u = -f''(\theta) \sin \theta + f'(\theta) \cos \theta$ and $v = f''(\theta) \cos \theta + f'(\theta) \sin \theta$, then

$$\int \left[\left(\frac{du}{d\theta} \right)^2 + \left(\frac{dv}{d\theta} \right)^2 \right]^{\frac{1}{2}} d\theta =$$

- (A) $f(\theta) - f''(\theta) + C$
(B) $f(\theta) + f''(\theta) + C$
(C) $f'(\theta) + f''(\theta) + C$
(D) $f'(\theta) - f''(\theta) + C$
(E) $f(\theta) + f'(\theta) + C$

Space for rough work

107. $\int \frac{e^{6\log_e x} - e^{5\log_e x}}{e^{4\log_e x} - e^{3\log_e x}} dx$ is equal to

(A) $\frac{x^3}{3} + C$ (B) $\frac{x^2}{2} + C$ (C) $\frac{x^2}{3} + C$ (D) $\frac{-x^3}{3} + C$ (E) $x + C$

108. $\int e^x \left(\frac{1-x}{1+x^2} \right)^2 dx$ is equal to

(A) $e^x \left(\frac{1-x}{1+x^2} \right) + C$ (B) $e^x \left(\frac{1}{1+x^2} \right) + C$ (C) $e^x \left(\frac{1+x}{1+x^2} \right) + C$

(D) $e^x \left(\frac{1-x}{(1+x^2)^2} \right) + C$ (E) $e^x \left(\frac{1}{(1+x^2)^2} \right) + C$

109. $\int \frac{x^4 - 1}{x^2(x^4 + x^2 + 1)^{\frac{1}{2}}} dx$ is equal to

(A) $\sqrt{\frac{x^4 + x^2 + 1}{x}} + C$ (B) $\frac{x^2}{\sqrt{x^4 + x^2 + 1}} + C$ (C) $x(x^4 + x^2 + 1)^{\frac{3}{2}} + C$

(D) $\frac{\sqrt{x^4 + x^2 + 1}}{x} + C$ (E) $\sqrt{x^4 + x^2 + 1} + C$

Space for rough work

110. $\int \frac{\cos x - \sin x}{1 + 2 \sin x \cos x} dx$ is equal to

- (A) $-\frac{1}{\cos x - \sin x} + C$ (B) $\frac{\cos x + \sin x}{\cos x - \sin x} + C$ (C) $-\frac{1}{\sin x + \cos x} + C$
(D) $\frac{x}{\sin x + \cos x} + C$ (E) $\tan x \sec x + C$

111. $\int \frac{1}{x} (\log_{ex} e) dx$ is equal to

- (A) $\log_e(1 - \log_e x) + C$ (B) $\log_e(\log_e ex - 1) + C$ (C) $\log_e(\log_e x - 1) + C$
(D) $\log_e(\log_e x + x) + C$ (E) $\log_e(1 + \log_e x) + C$

112. The value of $\int_1^e 10^{\log_e x} dx$ is equal to


- (A) $10 \log_e(10e)$ (B) $\frac{10e-1}{\log_e 10e}$ (C) $\frac{10e}{\log_e 10e}$
(D) $(10e) \log_e(10e)$ (E) $\frac{10}{\log_e(10e)}$

Space for rough work

113. The area between the curve $y = 1 - |x|$ and the x -axis is equal to
 (A) 1 sq.unit (B) $\frac{1}{2}$ sq.unit (C) $\frac{1}{3}$ sq.unit (D) 2 sq.units (E) 3 sq.units

114. The value of $\int_{e^{-1}}^e \frac{dt}{t(1+t)}$ is equal to
 (A) 0 (B) $\log\left(\frac{e}{1+e}\right)$ (C) $\log\left(\frac{1}{1+e}\right)$
 (D) $\log(1+e)$ (E) 1

115. The figure shows a triangle AOB and the parabola $y = x^2$.
 The ratio of the area of the triangle AOB to the area
 of the region AOB of the parabola $y = x^2$ is equal to



- (A) $\frac{3}{5}$ (B) $\frac{3}{4}$ (C) $\frac{7}{8}$ (D) $\frac{5}{6}$ (E) $\frac{2}{3}$

Space for rough work

116. The value of $\int_{-2}^4 |x+1| dx$ is equal to
(A) 12 (B) 14 (C) 13 (D) 16 (E) 15

117. The solution of $\cos y \frac{dy}{dx} = e^{x+\sin y} + x^2 e^{\sin y}$ is
(A) $e^x - e^{-\sin y} + \frac{x^3}{3} = C$ (B) $e^{-x} - e^{-\sin y} + \frac{x^3}{3} = C$ (C) $e^x + e^{-\sin y} + \frac{x^3}{3} = C$
(D) $e^x - e^{\sin y} - \frac{x^3}{3} = C$ (E) $e^x - e^{\sin y} + \frac{x^3}{3} = C$

118. The order and degree of the differential equation $\left(1 + \left(\frac{dy}{dx}\right)^2\right)^{\frac{3}{4}} = \left(\frac{d^2y}{dx^2}\right)^{\frac{1}{3}}$ is
(A) (2, 4) (B) (2, 3) (C) (6, 4) (D) (6, 9) (E) (2, 12)

Space for rough work

Mathrubhumi
Education

119. The integrating factor of the differential equation $(y \log y)dx = (\log y - x)dy$ is
- (A) $\frac{1}{\log y}$ (B) $\log(\log y)$ (C) $1 + \log y$ (D) $\frac{1}{\log(\log y)}$ (E) $\log y$
120. The solution of the differential equation $\frac{dy}{dx} = \frac{1}{x+y^2}$ is
- (A) $y = -x^2 - 2x - 2 + ce^x$ (B) $y = x^2 + 2x + 2 - ce^x$
(C) $x = -y^2 - 2y + 2 - ce^y$ (D) $x = -y^2 - 2y - 2 + ce^y$
(E) $x = y^2 + 2y + 2 - ce^y$

Space for rough work

Mathrubhumi
Education

Space for rough work

**Mathrubhumi
Education**

**Mathrubhumi
Education**

Maths-09/B1

31

Mathrubhumi Education

BLANK PAGE

Mathrubhumi Education